



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2012

Selection for tetracycline-resistant *Chlamydia suis* in treated pigs

Borel, Nicole ; Regenscheit, N ; Di Francesco, A ; Donati, M ; Markov, J ; Masserey, Y ; Pospischil, A

Abstract: The aim of this study was to investigate *Chlamydia suis* in a pig farm with an outbreak of conjunctivitis and diarrhea. Eye swabs and pooled fecal samples were investigated for the presence of *C. suis* by real-time PCR and ArrayTube microarray. Samples positive for *C. suis* by ArrayTube microarray assay were further tested for the presence of the tet(C) resistance gene by PCR. In the first examination, *C. suis* was identified in 12 six-week-old pigs showing conjunctivitis. Of these, the tet(C) gene-coding region was amplified in one pooled fecal sample and one eye swab, respectively. After oral treatment with tetracycline, clinical symptoms disappeared. Subsequently, all eye swabs investigated from 10 healthy pigs were positive for *C. suis* and the tet(C) gene-coding region. The present study reports rapid selection for tetracycline-resistant *C. suis* after antibiotic treatment.

DOI: <https://doi.org/10.1016/j.vetmic.2011.10.011>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-54309>

Journal Article

Accepted Version

Originally published at:

Borel, Nicole; Regenscheit, N; Di Francesco, A; Donati, M; Markov, J; Masserey, Y; Pospischil, A (2012). Selection for tetracycline-resistant *Chlamydia suis* in treated pigs. *Veterinary Microbiology*, 156(1-2):143-146.

DOI: <https://doi.org/10.1016/j.vetmic.2011.10.011>

Accepted Manuscript

Title: Selection for Tetracycline-Resistant *Chlamydia suis* in Treated Pigs

Authors: Nicole Borel, Nadine Regenscheit, Antonietta Di Francesco, Manuela Donati, Jenny Markov, Yvonne Masserey, Andreas Pospischil



PII: S0378-1135(11)00554-2
DOI: doi:10.1016/j.vetmic.2011.10.011
Reference: VETMIC 5497

To appear in: *VETMIC*

Received date: 25-7-2011
Revised date: 4-10-2011
Accepted date: 7-10-2011

Please cite this article as: Borel, N., Regenscheit, N., Di Francesco, A., Donati, M., Markov, J., Masserey, Y., Pospischil, A., Selection for Tetracycline-Resistant *Chlamydia suis* in Treated Pigs, *Veterinary Microbiology* (2010), doi:10.1016/j.vetmic.2011.10.011

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Selection for Tetracycline-Resistant *Chlamydia suis* in Treated Pigs

Nicole Borel^{1a}, Nadine Regenscheit^{1a}, Antonietta Di Francesco², Manuela Donati³,
Jenny Markov⁴, Yvonne Masserey⁴, Andreas Pospischil¹

1 Institute for Veterinary Pathology, University of Zurich, Vetsuisse Faculty,
Winterthurerstrasse 268, CH-8057 Zurich, Switzerland

2 Department of Veterinary Medical Sciences, University of Bologna, Via Tolara di
Sopra, 50, 40064 Ozzano Emilia (BO), Italy

3 Section of Microbiology DESOS, Policlinico S. Orsola, University of Bologna, Via
Massarenti 9, 40138 Bologna, Italy

4 SUISAG pig health service (PHS), Winterthurerstrasse 260, CH-8057 Zurich,
Switzerland

^aThese two authors contributed equally to this study

Corresponding author:

Nicole Borel, DVM, FVH, Dipl. ECVP
Institute of Veterinary Pathology, Vetsuisse Faculty
University of Zurich, Winterthurerstrasse 268
CH-8057 Zurich, Switzerland
Tel.: +41-44-635-8576
Fax.: +41-44-635-8934
Email: n.borel@access.uzh.ch

Abstract

The aim of this study was to investigate *Chlamydia suis* in a pig farm with an outbreak of conjunctivitis and diarrhea. Eye swabs and pooled fecal samples were investigated for the presence of *C. suis* by real-time PCR and ArrayTube microarray. Samples positive for *C. suis* by ArrayTube microarray assay were further tested for the presence of the *tet*(C) resistance gene by PCR. In the first examination, *C. suis* was identified in 12 six-week old pigs showing conjunctivitis. Of these, the *tet*(C) gene-coding region was amplified in one pooled fecal sample and one eye swab, respectively. After oral treatment with Tetracycline, clinical symptoms disappeared. Subsequently, all eye swabs investigated from ten healthy pigs were positive for *C. suis* and the *tet*(C) gene-coding region. The present study reports rapid selection for Tetracycline-resistant *C. suis* after antibiotic treatment.

Key words

Chlamydia suis, conjunctivitis, diarrhea, selection, swine, Tetracycline-resistance

Introduction

Chlamydia (*C.*) *suis* is associated with respiratory disease, diarrhea and conjunctivitis in pigs (Pospischil et al., 2010). Chlamydiae are commonly found in the intestine of pigs mostly associated with subclinical infections (Nietfeld et al., 1997). However, *C. suis* caused enteritis when inoculated in gnotobiotic piglets (Rogers and Andersen, 1996; Guscetti et al., 2009) or caused histologically intestinal lesions in asymptomatic young weanling pigs (Rogers and Andresen, 2000). Tetracycline-resistant *C. suis* strains have been described in the US (Lenart et al., 2001) and in Italy (Di Francesco et al., 2008). In vitro, chlamydial inclusions contain aberrant forms when cultured in addition of tetracycline indicating the induction of persistence by the addition of

antibiotics (Lenart et al., 2001). In vivo, aberrant chlamydial developmental forms in the gastrointestinal tract of pigs spontaneously and experimentally infected with *C. suis* have been shown (Pospischil et al., 2009). In the latter study, group A consisted of conventionally reared pigs from a farm in the US and was used for experimental infection with *Salmonella typhimurium*. These pigs were by chance detected to be naturally infected with *C. suis*, but unavailability of fresh tissue material prevented the amplification of the Tetracycline (*tet*(C)) resistance gene (data not shown). Group B consisted of gnotobiotic piglets experimentally infected with the *C. suis* strain S45/6 which is known to lack the *tet*(C) gene-coding region. Interestingly, this strain was harvested originally in the late 1960s from feces of an asymptomatic pig in Austria indicating the presence of non-resistant *C. suis* strains in Europe at that time. Tetracyclines are widely used in veterinary medicine because they are relatively inexpensive and have a broadspectrum activity. To date, selection of Tetracycline-resistance in *C. suis* in pigs after antibiotic treatment is not reported.

Material and Methods

In April 2010, a Swiss breeding pig farm consisting of approximately 400 animals had problems of wasting, decreased body weight, diarrhea and conjunctivitis in weaned pigs. Fertility parameters in breeding sows were normal. A vaccination protocol for *Lawsonia intracellularis*, porcine Circovirus, porcine Parvovirus, *Escherichia coli*, *Clostridium* sp. and Erysipelas was performed in the herd. Deworming and iron prophylaxis was regularly followed and the hygiene management of the herd was good. Eye swabs of 12 six-week old pigs showing conjunctivitis (conjunctival chemosis and reddening, ocular discharge) and pooled fecal samples (n=3) of their housing compartments were taken for further examination. Subsequently, all pigs of the farm were treated orally with tetracycline for 3 weeks. In August 2010, no clinical

symptoms were recorded in the farm. Control eye swabs were taken from ten healthy pigs.

DNA of all samples (eye swabs: n = 22; pooled fecal samples: n = 3) was extracted using the DNeasy Blood and Tissue Kit (QIAGEN, Hilden, Germany) according to the manufacturer's instructions. All samples were examined on an ABI 7500 instrument (Applied Biosystems, Foster City, CA, USA) using the 23S rRNA gene-based *Chlamydiaceae* family-specific real-time PCR as described previously (Ehrlich et al., 2006). Briefly, this method includes primers Ch23S-F (5'-CTGAAACCAGTAGCTTATAAG CGGT-3'), Ch23S-R (5'-ACCTCGCCGTTTAACTTAACTCC-3'), and probe Ch23S-p (FAM-CTCATCATGCAAAAGGCACGCCG-TAMRA) and an internal amplification control consisting of primers EGFP-1-F (5'-GACCACTACCAGCAGAACAC-3'), EGFP-10-R (3'-CTTGTACAGCTCGTCCATGC-5') and probe EGFP-HEX (HEX-AGCACCCAGTCCGCCCTGAGCA-BHQ1). A 111-bp product specific for members of the family *Chlamydiaceae* is produced as well as a 177-bp product for the internal amplification control. A cycle threshold (Ct value) of < 38.00 was considered as positive, and all samples were tested at least in duplicate. The samples with positive Ct values were examined using the species-specific 23S ArrayTube (AT) microarray (Alere, Jena, Germany) assay (Borel et al., 2008). Samples positive for *C. suis* by AT microarray assay were further tested for the presence of the *tet(C)* resistance gene by a PCR assay amplifying a 525 base pair product of the *tet(C)* gene-coding region. The reaction was performed according to Dugan and others (2004), using following forward and reverse oligonucleotide primers: CS43 5'-AGCACTGTCCGACCGCTTTG-3' and CS47 5'-TCCTCGCCGAAAATGACCC-3'. The S45 tetracycline-sensitive and the MS08 tetracycline-resistant *C. suis* strains were used as negative and positive control, respectively. The amplicons were

purified using a commercially available kit (QIAquick PCR Purification Kit, QIAGEN, Hilden, Germany), sequenced (Bio-Fab Research, Rome, Italy) and identified through BLAST search (www.ncbi.nlm.nih.gov/blast/).

Results and Discussion

Real-time PCR for *Chlamydiaceae* was positive in all 12 eye swabs and the pooled fecal samples (n=3) in April 2010. By AT microarray, *C. suis* was identified in all samples (Figure 1). Of these, the *tet(C)* gene-coding region was amplified in one pooled fecal sample and one eye swab, respectively. Amplicons carried an identical nucleotide sequence that showed 100% homology with those of the structural gene *tet(C)* described by Dugan and others (2004) (Gene Bank Accession AY428551).

Other infective agents causing conjunctivitis and diarrhea were excluded by laboratory investigations (data not shown). In August 2010, no clinical symptoms were recorded in the farm. All eye swabs (n = 10) collected from healthy pigs were positive for *C. suis* and the *tet(C)* gene-coding region.

Chlamydiae were reported as a cause of conjunctivitis in pigs in Nebraska in 1993 (Rogers et al., 1993). As in this study, weanling pigs were affected and they responded well to oxytetracycline treatment. At that time, no clear classification of the involved chlamydial species was performed. *C. suis* was found in a study in pigs with conjunctivitis but also in healthy controls from Switzerland and Germany (Becker et al., 2007). A concurrent outbreak of chlamydial disease and postweaning multisystemic wasting syndrome was recently described in weaned piglets in a pig production farm in Estonia (Schautteet et al., 2010). *C. suis* DNA was detected in eye swabs from boars, sows and gilts with conjunctivitis and also in feces of boars and sows. Therapy with doxycycline was not successful and clinical signs persisted. The authors allocated this to either to short treatment period, to prolonged shipment of the

1 samples, or to the presence of Tetracycline-resistant strains. In both studies (Becker
 2 et al., 2007; Schautteet et al., 2010), *C. suis*-strains were not further investigated for
 3 the Tetracycline-resistance gene. In another study (Reinhold et al., 2011), short-term
 4 treatment with enrofloxacin or enrofloxacin plus tiamulin was not effective in
 5 eliminating subclinical infections with *C. suis* in pigs. In the present study, clinical
 6 symptoms disappeared after antibiotic treatment but *C. suis* was not eliminated and
 7 strains harboring the *tet(C)* resistance gene were positively selected. As recently
 8 reviewed by Sandoz and Rockey (2011), the discovery of the *tet(C)* islands
 9 represents the first identification of antibiotic resistance acquired through horizontal
 10 gene transfer in any obligate intracellular bacteria. The mechanism of acquisition of
 11 the *tet(C)* islands in *C. suis* remains speculative. However, acquisition of DNA from
 12 other bacteria by *Chlamydia* that commonly infect the porcine intestinal tract such as
 13 *C. suis* is conceivable. The rapid selection for Tetracycline-resistant *C. suis* strains
 14 after antibiotic treatment in this study was surprising and possibly facilitated by the
 15 close contact of pigs in a relatively small farm (400 animals). In vitro experiments
 16 have demonstrated the acquisition of Tetracycline resistance by horizontal gene
 17 transfer from *C. suis* to clinical strains of *C. trachomatis* (Suchland et al., 2009). In
 18 vivo, such events are of particular public health concern if the *tet(C)* resistance gene
 19 is transferred from porcine chlamydial strains to human pathogens.
 20 *C. suis* was found in pooled fecal samples of diarrheic pigs in the present study.
 21 Prophylactic vaccination against common enteropathogenic agents was performed in
 22 this herd and other infectious agents were excluded by laboratory investigations
 23 (data not shown). This implies *C. suis* as the sole agent causing diarrhea in this herd,
 24 although mostly subclinical infections have been reported in previous field studies
 25 (Nietfeld et al., 1997). It is known that different isolates of *C. suis* exist with a high

degree of genetic diversity possibly leading to variation in virulence. Further characterization of these chlamydial strains is currently under investigation.

Conclusion

This study resulted in the first description of Tetracycline-resistant *C. suis* in a Swiss pig farm with conjunctivitis and diarrhea. Excretion of these strains by ocular and fecal route was present before and after antibiotic treatment. After tetracycline treatment, strains harboring the *tet(C)* gene-coding region were predominantly prevalent indicating selection of these under treatment conditions. The adaptive availability to acquire the *tet(C)*-resistant gene when exposed to selective pressure could have far-reaching consequences for the emergence of antibiotic resistance in chlamydiae and could pose a potential threat for the food safety. Further investigations in other European countries to detect *tet(C)*-resistant strains are needed to increase awareness and caution when treating pigs with antibiotics.

Conflict of interest statement

All authors declare there is no financial or personal relationships with other people or organizations that could have inappropriately influenced their work.

Acknowledgments

We are grateful to Carmen Kaiser of the laboratory of the Institute of Veterinary Pathology, University of Zurich for technical help.

References

Becker, A., Lutz-Wohlgroth, L., Brugnera, E., Lu, Z.H., Zimmermann, D.R., Grimm, F., Grosse Beilage, E., Kaps, S., Spiess, B., Pospischil, A., Vaughan, L., 2007.

- Intensively kept pigs pre-disposed to chlamydial associated conjunctivitis. J. Vet. Med. A. Physiol. Pathol. Clin. Med. 54, 307-313.
- Borel, N., Kempf, E., Hotzel, H., Schubert, E., Torgerson, P., Slickers, P., Ehricht, R., Tasara, T., Pospischil, A., Sachse, K., 2008. Direct identification of chlamydiae from clinical samples using a DNA microarray assay - A validation study. Mol. Cell. Probes. 22, 55-64.
- Di Francesco, A., Donati, M., Rossi, M., Pignanelli, S., Shurdhi, A., Baldelli, R., Cevenini, R., 2008. Tetracycline-resistant *Chlamydia suis* isolates in Italy. Vet. Rec. 163, 251-252.
- Dugan, J., Rockey, D.D., Jones, L., Andersen, A.A., 2004. Tetracycline resistance in *Chlamydia suis* mediated by genomic islands inserted into the chlamydial inv-like gene. Antimicrob. Agents. Chemother. 48, 3989-3995.
- Ehricht, R., Slickers, P., Goellner, S., Hotzel, H., Sachse, K., 2006. Optimized DNA microarray assay allows detection and genotyping of single PCR-amplifiable target copies. Mol. Cell. Probes. 20, 60-63.
- Guscetti, F., Schiller, I., Sydler, T., Heinen, E., Pospischil, A., 2009. Experimental enteric infection of gnotobiotic piglets with *Chlamydia suis* strain S45. Vet. Microbiol. 135, 157-168.
- Lenart, J., Andersen, A.A., Rockey, D.D., 2001. Growth and development of Tetracycline-resistant *Chlamydia suis*. Antimicrob. Agents. Chemother. 45, 2198-

2203.

Nietfeld, J.C., Leslie-Steen, P.L., Zeman, D.H., Nelson, D., 1997. Prevalence of intestinal chlamydial infection in pigs in the midwest, as determined by immunoperoxidase staining. *Am. J. Vet. Res.* 58, 260-264.

Pospischil, A., Borel, N., Andersen, A.A., 2010. *Chlamydia*. In: Gyles, C.L., Prescott, J.F., Songer, J.G., Thoen, C.O. (Eds.), *Pathogenesis of bacterial infections in animals*. Blackwell Publishing, pp. 575-87.

Pospischil, A., Borel, N., Chowdhury, E.H., Guscetti, F., 2009. Aberrant chlamydial developmental forms in the gastrointestinal tracts of pigs spontaneously and experimentally infected with *Chlamydia suis*. *Vet. Microbiol.* 135, 147-56.

Reinhold, P., Liebler-Tenorio, E., Sattler, S., Sachse, K., 2011. Recurrence of *Chlamydia suis* infection in pigs after short-term antimicrobial treatment. *Vet. J.* 187, 405-407.

Rogers, D.G., Andersen, A.A., 1996. Intestinal lesions caused by two swine chlamydial isolates in gnotobiotic piglets. *J. Vet. Diagn. Invest.* 8, 433-440.

Rogers, D.G., Andersen, A.A., 2000. Intestinal lesions caused by a strain of *Chlamydia suis* in weanling pigs infected at 21 days of age. *J. Vet. Diagn. Invest.* 12, 233-239.

- Rogers, D.G., Andersen, A.A., Hogg, A., Nielsen, D.L., Huebert, M.A., 1993. Conjunctivitis and keratoconjunctivitis associated with chlamydiae in swine. J. Am. Vet. Med. Assoc. 203, 1321-1323.
- Sandoz, K.M., Rockey, D.D., 2010. Antibiotic resistance in Chlamydiae. Future Microbiol. 5, 1427-1442.
- Schautteet, K., Beeckman, D.S.A., Delava, P., Vanrompay, D., 2010. Possible pathogenic interplay between *Chlamydia suis*, *Chlamydophila abortus* and PCV-2 on a pig production farm. Vet Rec. 166, 329-333.
- Suchland, R.J., Sandoz, K.M., Jeffrey, B.M., Stamm, W.E., Rockey, D.D., 2009. Horizontal transfer of Tetracycline resistance among *Chlamydia* spp. in vitro. Antimicrob. Agents. Chemother. 53, 4604-4611.

Figure captions

Fig. 1. DNA-based ArrayTube Microarray species identification assay of eye swab from a pig positive for *Chlamydiaceae* by real-time PCR (mean Ct value 24.77). Barplot showing specific signals for genus *Chlamydia* (1) and species *Chlamydia suis* (2). Red bars represent the perfect-match probe signal, blue bars show the hybridization signal with the probe carrying one or two mismatches at the 3'-end.

